

ASYMPTOTIC BOUNDS FOR NORI'S CONNECTIVITY THEOREM

ANIA OTWINOWSKA

ABSTRACT. Let Y be a smooth complex projective variety. I study the cohomology of smooth families of hypersurfaces $\mathcal{X} \rightarrow B$ for $B \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ a codimension c subvariety. I give an asymptotically optimal bound on c and k when $d \rightarrow \infty$ for the space $H^k(Y \times B, \mathcal{X}, \mathbb{Q})$ to vanish, thus extending the validity of Lefschetz Hyperplane section Theorem and Nori's Connectivity Theorem [No]. Next, I construct in the limit case explicit families of higher Chow groups whose class does not vanish in $H^k(Y \times B, \mathcal{X}, \mathbb{Q})$. Some of them are indecomposable. This suggests that in the limit case the space $H^k(Y \times B, \mathcal{X}, \mathbb{Q})$ should be spanned by higher Chow groups, containing [No] and [O] as special cases (*cf.* conjecture 2).

0. INTRODUCTION

0.1. Nori's connectivity theorem. Let Y be a smooth complex proper algebraic variety of dimension $N + 1$ with an ample invertible sheaf $\mathcal{L}$ and $\pi: \mathcal{X} \rightarrow \mathbb{P}H^0(Y, \mathcal{L})$ the universal family of hypersurfaces of class $c_1(\mathcal{L})$. For any $b \in \mathbb{P}H^0(Y, \mathcal{L})$, let $X_b = \pi^{-1}(b)$ be the corresponding hypersurface of Y .

The Lefschetz hyperplane section theorem asserts that for $k + 1 \in \{0, \dots, N\}$,

$$H^{k+1}(Y, X_b, \mathbb{Z}) = 0.$$

Nori's connectivity theorem extends this result to generically defined cohomology classes of degree all the way up to $2N = \dim_{\mathbb{R}} X$. More precisely, let T be a smooth variety and $\phi: T \rightarrow \mathbb{P}H^0(Y, \mathcal{L})$ a differentiable map. Let $\mathcal{X}_T = \mathcal{X} \times_{\mathbb{P}H^0(Y, \mathcal{L})} T$ be the universal hypersurface over T and let $\mathcal{Y}_T = Y \times T$. By the Leray spectral sequence, the Lefschetz Hyperplane Section Theorem implies that for $k + 1 \in \{0, \dots, N\}$,

$$H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Z}) = 0.$$

With stronger assumptions one can say more. Assume that T is a smooth algebraic variety and that $\phi: T \rightarrow \mathbb{P}H^0(Y, \mathcal{L})$ is a smooth (hence dominant) morphism. Let $\mathcal{O}(1)$ be a fixed very ample invertible sheaf on Y and assume $\mathcal{L} = \mathcal{O}(d)$ for some positive integer d .

Theorem (Nori's connectivity theorem [No]). *For d large enough and $k+1 \in \{0, \dots, 2N\}$,*

$$H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q}) = 0.$$

Note that Nori's connectivity theorem holds more generally for families of complete intersections in Y (instead of only hypersurfaces). The more general setting is motivated by the construction of algebraic cycles which are homologically but not algebraically equivalent to zero. Since in this paper I do not consider applications to algebraic equivalence,

I restrict myself to the case of hypersurfaces, which involves all the ideas and simplifies notations.

Despite its topological nature, the proof of Nori's connectivity theorem relies on Hodge theory (cf. [No], section 1). Namely, let $\Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p$ be the coherent sheaf of $\mathcal{O}_{Y_T}$ -modules defined by the short exact sequence

$$0 \rightarrow \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p \rightarrow \Omega_{\mathcal{Y}_T}^p \rightarrow j_{T*} \Omega_{\mathcal{X}_T}^p \rightarrow 0,$$

where $j_T: \mathcal{X}_T \rightarrow \mathcal{Y}_T$ is the closed immersion. Then $H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = \mathbb{H}^{p+q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^\bullet)$. Following Nori, let

$$G^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = \text{Im}(\mathbb{H}^{p+q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^{\bullet \geq p}) \rightarrow H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}))$$

denote the Hodge-Grothendick filtration. There is a natural inclusion

$$F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) \subset G^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}),$$

where $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$ denotes the Hodge filtration of the mixed Hodge structure on $H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T)$. Hence if for some positive integer k one has $H^{q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p) = 0$ for all $p + q + 1 \leq k + 1$ and $q \leq (k + 1)/2$, then by the Hodge–Fröhlicher spectral sequence $G^{k+1-[(k+1)/2]} H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$, hence $F^{k+1-[(k+1)/2]} H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ and a mixed Hodge structures argument implies $H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q}) = 0$. Hence the connectivity theorem follows from the following more precise Hodge-theoretical statement

Theorem (Nori). *For d large enough, $p + q + 1 \leq 2N$ and $q + 1 \leq N$,*

$$H^{q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p) = 0.$$

0.2. Vanishing theorems. The first goal of this paper is to investigate the minimal assumptions for Nori's connectivity theorem to extend. Natural questions are:

- how large does d need to be?
- what happens if the map $\phi: T \rightarrow \mathbb{P}H^0(Y, \mathcal{O}(d))$ fails to be dominant, i.e if one considers families of hypersurfaces which are not general, but close to being general?
- can the above assertions be made more precise for a fixed integer k (in Nori's connectivity theorem), and for fixed integers p and q (in the Hodge-theoretical version)?

The first question was studied in Nagel [N] and Voisin [V2]. In this paper I give the following asymptotic answer to the above questions.

Let T be a smooth algebraic variety with an algebraic morphism $\phi: T \rightarrow \mathbb{P}H^0(Y, \mathcal{O}(d))$, such that $\mathcal{X}_T$ is smooth. I assume that for all $t \in T$ the cokernel of the differential map $d\phi(t): T_{T,t} \rightarrow T_{\mathbb{P}H^0(Y, \mathcal{O}(d)), \phi(t)}$ is of dimension at most an integer c .

Theorem 1. *For all $\varepsilon \in]0, 1[$ and for all $p \in \mathbb{N}$ and for $q + 1 \in \{0, \dots, N\}$, there is an integer D depending only on ε, p, Y and $\mathcal{O}(1)$, such that for all $d \geq D$,*

$$H^{q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p) = 0$$

provided that

$$c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}.$$

A straightforward adaptation of Nori's argument sketched above shows that in the numerical situation of Theorem 1 $G^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$, hence $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$, and the following topological statement holds.

Corollary . *For all $\varepsilon \in]0, 1[$ and $k + 1 \in \{0, \dots, 2N\}$ there is an integer D depending only on ε , Y and $\mathcal{O}(1)$ such that for all $d \geq D$,*

$$H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q}) = 0$$

provided that

$$c \leq (1 - \varepsilon) \frac{d^{N-[k/2]}}{(N - [k/2])!}.$$

The following remarks show that the bounds given by Theorem 1 and the Corollary are close to being asymptotically optimal:

0.2.1. *Case $k + 1 \in \{0, \dots, N\}$.* By the Lefschetz hyperplane section theorem and the Leray spectral sequence one has $H^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q}) = 0$ with no assumption on c . (similarly, by the holomorphic Leray spectral sequence one has $H^{q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p) = 0$ for $p + q < N$ with no assumption on c).

0.2.2. *Case $k + 1 \in \{N + 1, \dots, 2N\}$.* I show in the next section (see Theorem 3 below) that under a weak assumption on Y , the bound on c is asymptotically optimal for any couple of fixed integers p and q such that $N/2 \leq q \leq N - 1$ and $p \geq q$. This implies that the bound given by the Corollary is asymptotically optimal.

0.2.3. *Case $k + 1 \geq 2N + 1$.* One has $H^{2N+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q}) \neq 0$ for any positive integer c , $d > N$ and $T = \mathcal{X}_B$, where $B \subset \mathbb{P}H^0(Y, \mathcal{O}(1))$ is any smooth codimension c subvariety contained in the locus of smooth hypersurfaces. Indeed, the condition $d > N$ implies $H^{0,N}(X_b, \mathbb{Q}) \neq 0$ for all $b \in B$, hence (cf. [No], section 0.2) the cohomology class of the diagonal embedding of $\mathcal{X}_B$ in $\mathcal{X}_T = \mathcal{X}_B \times_B \mathcal{X}_B$ is an element of $H^{2N}(\mathcal{X}_T, \mathbb{Q})$ which does not belong to $j_T^* H^{2N}(\mathcal{Y}_T, \mathbb{Q})$, hence defines a non-zero element of $H^{2N+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{Q})$.

0.3. **Some examples of algebraic classes.** The aim of this section is to give explicit constructions of *algebraic* classes in $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$ for $q + 1 \in \{0, \dots, N\}$, $p \geq q$ and c close to the bound given by theorem 1.

0.3.1. *The regulator map.* For any smooth algebraic variety X and positive integers $p \geq q$ the higher Chow groups $CH^p(X, p - q)$ are defined in [B]. In this paper they are always implicitly assumed to be with $\mathbb{Q}$ coefficients. They come with the Deligne class map

$$cl_{p,p-q}(\mathcal{D}): CH^p(X, p - q) \rightarrow H_{\mathcal{D}}^{p+q}(X, \mathbb{Q}(p)),$$

where $H_{\mathcal{D}}^{p+q}(X, \mathbb{Q}(p))$ denotes Deligne cohomology. There is a natural map

$$r_{p,p-q}: H_{\mathcal{D}}^{p+q}(X, \mathbb{Q}(p)) \rightarrow F^p H^{p+q}(X, \mathbb{C}),$$

where $F^p\mathrm{H}^{p+q}(X, \mathbb{C})$ denotes the Hodge filtration. Composing $\mathrm{cl}_{p,p-q}(\mathcal{D})$ with $r_{p,p-q}$ one gets the regulator map

$$\mathrm{reg}_{p,p-q}: \mathrm{CH}^p(X, p-q) \rightarrow F^p\mathrm{H}^{p+q}(X, \mathbb{C}).$$

Note that this map is identically zero for $p > q$ whenever X is proper.

Consider the case $X = \mathcal{X}_T$ (note that $\mathcal{X}_T$ is *not* proper). Let

$$\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v = \mathrm{coker}(j_T^*: \mathrm{H}^{p+q}(\mathcal{Y}_T, \mathbb{C}) \rightarrow \mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C}))$$

denote the *vanishing* cohomology, let $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v = F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C}) \cap \mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ and let

$$\overline{\mathrm{reg}}_{p,p-q}: \mathrm{CH}^p(\mathcal{X}_T, p-q) \rightarrow F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v \subset F^p\mathrm{H}^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$$

be the map deduced from $\mathrm{reg}_{p,p-q}$ by projection on $\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$.

An element of $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ (or $F^p\mathrm{H}^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$) will be called *algebraic* if it lies in the image of $\overline{\mathrm{reg}}_{p,p-q}$.

0.3.2. Chow groups. In the case $p = q$ higher Chow groups are Chow groups and the regulator is the class map. I then write $\mathrm{CH}^p(\mathcal{X}_T)$ for $\mathrm{CH}^p(\mathcal{X}_T, 0)$, cl_p for $\mathrm{reg}_{p,0}$ and $\overline{\mathrm{cl}}_p$ for $\overline{\mathrm{reg}}_{p,0}$.

The following is a variation of my result in [O].

Theorem 2. *Let p, q, b and c be positive integers such that $p + q = N$, $q \in \{[(N+1)/2], \dots, N-1\}$, $b \geq 1$ and $c \leq b \frac{d^{N-q}}{(N-q)!}$. Assume $\phi(T) \subset \mathbb{P}^0(Y, \mathcal{O}(d))$ lies in the locus parametrizing smooth hypersurfaces. Then there is an integer D depending only on b, Y and $\mathcal{O}(1)$ such that for all $d \geq D$ the following holds.*

If N is even and $p = q = N/2$ then the space $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ is spanned via $\overline{\mathrm{cl}}_p$ by classes of flat families of algebraic cycles in $\mathcal{X}_T$ of codimension p and degree at most b .

If $q > N/2$ then $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v = 0$.

0.3.3. Higher Chow groups. I now fulfill the promise made in section 0.2.2.

Theorem 3. *Let p and q be positive integers such that $p \geq q$ and $q \in \{[(N+1)/2], \dots, N-1\}$. Assume that $Y \subset \mathbb{P}^0(Y, \mathcal{O}(1))^\vee$ contains a codimension $q+1$ linear subvariety V and let $T \subset \mathbb{P}^0(Y, \mathcal{O}(d))$ be the subvariety parametrising smooth hypersurfaces containing V . Then for $d \gg 0$ the variety T is of codimension $c \sim_{d \rightarrow \infty} \frac{d^{N-q}}{(N-q)!}$, and there is a higher cycle $Z \in \mathrm{CH}^p(\mathcal{X}_T, p-q)$ such that $\overline{\mathrm{reg}}_{p,p-q}(Z) \neq 0$.*

Hence in this situation the bound on c in Theorem 1 and its corollary are asymptotically optimal.

Note that the assumption on Y holds for $Y = \mathbb{P}_{\mathbb{C}}^{N+1}$, which should be regarded as the main example.

0.3.4. Indecomposable Higher Chow groups. For all $i \in \{0, \dots, p\}$ and $j \in \{0, \dots, p-q\}$ there are natural product maps

$$\psi_{i,j}: \mathrm{CH}^{p-i}(\mathcal{X}_T, p-q-j) \otimes \mathrm{CH}^i(\mathcal{X}_T, j) \rightarrow \mathrm{CH}^p(\mathcal{X}_T, p-q).$$

I say that a higher cycle in $\mathrm{CH}^p(\mathcal{X}_T, p - q)$ is *decomposable* if it belongs to the subgroup spanned by the images of the maps $\psi_{i,j}$ for $(i, j) \notin \{(0, 0), (p, p - q)\}$, and that it is *indecomposable* otherwise.

The proof of Theorem 3 relies on an explicit construction of higher cycles in $\mathrm{CH}^p(\mathcal{X}_T, p - q)$. More precisely, since $\mathrm{CH}^1(\mathcal{X}_T, 1) \simeq \mathrm{H}^0(\mathcal{X}_T, \mathcal{O}_{\mathcal{X}_T}^*) \simeq \mathrm{H}^0(T, \mathcal{O}_T^*)$, iterating the above map $p - q$ times I obtain the map

$$\psi_{1,1}^{p-q}: \mathrm{CH}^q(\mathcal{X}_T) \otimes \mathrm{H}^0(T, \mathcal{O}_T^*)^{p-q} \rightarrow \mathrm{CH}^p(\mathcal{X}_T, p - q).$$

For $p > q$ the higher cycles in $\mathrm{CH}^p(\mathcal{X}_T, p - q)$ I construct are in the image of $\psi_{1,1}^{p-q}$. Since $\psi_{1,1}^{p-q}$ commutes to the restriction to a fiber, it follows that the restriction of these higher cycles to any fiber X_t , $t \in T$, is decomposable.

My result should be compared to Voisin's [V2], who studies the case $Y = \mathbb{P}_{\mathbb{C}}^{N+1}$, $p = N$, $q = N - 1$, $d = 2N$ and $T \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ an open subset (this implies $c = 0$). She constructs a family of higher cycles in $\mathrm{CH}^N(\mathcal{X}_T, 1)$ that limits the validity of Nori's connectivity theorem and whose restriction to a very general fiber X_t for $t \in T$ is indecomposable. This suggests that for all p, q such that $p + q \geq N$ and $p \geq q$ there might be higher cycles in $\mathrm{CH}^p(\mathcal{X}_T, p - q)$ for $c \sim_{d \rightarrow \infty} \frac{d^{N-q}}{(N-q)!}$ whose restriction to a very general fiber X_t , $t \in T$, is indecomposable. Supporting this intuition, I prove the following slightly weaker result for $Y = \mathbb{P}^{N+1}$, N even, $q = N/2$ and $p = q + 1$.

Theorem 4. *Assume N even, $q = N/2$ and that $Y \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(1))^\vee$ contains a codimension q linear subvariety. Then for $d \gg 0$ there is a subvariety $U \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ of codimension $c \sim_{d \rightarrow \infty} 2 \frac{d^{N-q}}{(N-q)!}$, and a higher cycle $Z_U \in \mathrm{CH}^{q+1}(\mathcal{X}_U, 1)$ such that*

- $\overline{\mathrm{reg}}_{q+1,1}(Z_U) \neq 0$, and
- the restriction of Z_U to a very general fiber X_u for $u \in U$ is indecomposable.

The cycle Z_U is constructed explicitly and the proof of indecomposability relies on Theorem 2.

I believe that Theorem 4 should generalize as follows.

Hope. *Let p and q be positive integers such that $p > q$ and $q \in \{[(N+1)/2], \dots, N-1\}$. Assume $Y = \mathbb{P}^{N+1}$. Then for $d \gg 0$ there is a subvariety $U \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ of codimension $c \sim_{d \rightarrow \infty} (p - q + 1) \frac{d^{N-q}}{(N-q)!}$ and a higher cycle $Z_U \in \mathrm{CH}^p(\mathcal{X}_U, p - q)$ such that*

- $\overline{\mathrm{reg}}_{p,p-q}(Z_U) \neq 0$, and
- the restriction of Z_U to a very general fiber X_u , $u \in U$, is indecomposable.

Theorem 4 is the case $p = N/2 + 1$, $q = N/2$.

Note that $c \sim_{d \rightarrow \infty} 2 \frac{d^{N-q}}{(N-q)!}$ in Theorem 4 (and $c \sim_{d \rightarrow \infty} (p - q + 1) \frac{d^{N-q}}{(N-q)!}$ in its conjectural generalisation) while Theorem 3 gives examples of decomposable cycles in $\mathrm{CH}^p(\mathcal{X}_T, p - q)$ with a non zero vanishing cohomology class for $c \sim_{d \rightarrow \infty} \frac{d^{N-q}}{(N-q)!}$. This suggests that the bound on c in Theorem 4 may not be optimal.

Question. *Is there an indecomposable higher cycle $Z_U \in \mathrm{CH}^p(\mathcal{X}_U, p - q)$ such that $\overline{\mathrm{reg}}_{p,p-q}(Z_U) \neq 0$ for $c \sim_{d \rightarrow \infty} \frac{d^{N-q}}{(N-q)!}$?*

0.4. Algebraicity conjecture. The results of Section 0.3 suggest that for c slightly bigger than the bound $\frac{d^{N-q}}{(N-q)!}$ given by Theorem 1 the space $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{Q})_v$ should be algebraic.

More precisely, I believe the following should hold

Conjecture 1. *For all integers $b \in \mathbb{N}^*$ and $p \in \mathbb{N}$ there is an integer D depending only on b , p , Y and on $\mathcal{O}(1)$, such that for all $d \geq D$ and $0 \leq q \leq N-1$, if*

$$c \leq b \frac{d^{N-q}}{(N-q)!}$$

then the map $\overline{\mathrm{reg}}_{p,p-q} : \mathrm{CH}^p(\mathcal{X}_T, p-q) \otimes \mathbb{C} \rightarrow F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ is surjective.

By convention, $\mathrm{CH}^p(\mathcal{X}_T, p-q) = 0$ for $p < q$.

By Theorem 2, Conjecture 1 holds for $p+q = N$, $q \geq N/2$ provided that $\phi(T) \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ lies in the locus parametrizing smooth hypersurfaces.

Mimicking Nori's argument, it is easy to see that Conjecture 1 implies the following purely topological statement.

Conjecture 2. *For every positive integer b there is an integer D depending only on b , Y and $\mathcal{O}(1)$ such that for all $d \geq D$ and $k \in \{N, \dots, 2N-1\}$ and for*

$$c \leq b \frac{d^{N-[k/2]}}{(N-[k/2])!},$$

- *if k is even then the map $\overline{\mathrm{cl}}_{k/2} : \mathrm{CH}^{k/2}(\mathcal{X}_T) \otimes \mathbb{C} \rightarrow F^{k/2}\mathrm{H}^k(\mathcal{X}_T, \mathbb{C})_v$ is surjective;*
- *if k is odd then the maps*

$$\overline{\mathrm{reg}}_{(k+1)/2,1} : \mathrm{CH}^{(k+1)/2}(\mathcal{X}_T, 1) \otimes \mathbb{C} \rightarrow \mathrm{H}^k(\mathcal{X}_T, \mathbb{C})_v, \quad \text{and}$$

$$\psi \circ \overline{\mathrm{reg}}_{(k+1)/2,1} : \mathrm{CH}^{(k+1)/2}(\overline{\mathcal{X}}_T, 1) \otimes \mathbb{C} \rightarrow \mathrm{H}^k(\overline{\mathcal{X}}_T, \mathbb{C})_v \rightarrow \mathrm{H}^k(\mathcal{X}_T, \mathbb{C})_v$$

generate $\mathrm{H}^{k+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})_v$.

Here $\overline{\mathcal{X}}_T$ is the variety isomorphic to $\mathcal{X}_T$ as a real manifold and endowed with the opposite complex structure and ψ is the geometric frobenius.


Note that for $p+q < N$ Conjecture 1 and for $k < N$ Conjecture 2 are trivially true since the target cohomology spaces vanish (cf. section 0.2.1).

For k even Theorem 1 implies that in the numerical situation of Conjecture 2, one has $F^{k/2+1}\mathrm{H}^k(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$, hence by a mixed Hodge structures argument the space $\mathrm{H}^k(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$ is of pure Hodge type $(k/2, k/2)$ (like in [No], section 0.2), hence the subspace $\mathrm{H}^k(\mathcal{X}_T, \mathbb{C})_v$ is a sub-Hodge structure of pure Hodge type $(k/2, k/2)$. Hence for k even Conjecture 2 is implied by Hodge Conjecture.

0.5. Summary. The figures summarise the results. Namely, for $c \leq b \frac{d^{N-q}}{(N-q)!}$

a : $F^p\mathrm{H}^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for any c by section 0.2.1.

b : $F^p\mathrm{H}^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq (1-\varepsilon) \frac{d^{N-q}}{(N-q)!}$ by Theorem 1 and $F^p\mathrm{H}^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ is spanned by classes of flat families of algebraic cycles of codimension $N/2$ and degree at most b for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Theorem 2, as predicted by Conjecture 1.



c : $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Theorem 2.

d : $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$ by Theorem 1 and one should still have $F^p H^{p+q}(\mathcal{X}_T, \mathbb{C})_v = 0$ for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Conjecture 1.

e : $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$ by Theorem 1 and $F^p H^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ should be spanned by $\overline{\text{cl}}_p(\text{CH}^p(\mathcal{X}_T))$ for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Conjecture 1; by Theorem 3 there is a variety Y such that $\overline{\text{cl}}_p(\text{CH}^p(\mathcal{X}_T)) \neq 0$ for $c \sim \frac{d^{N-q}}{(N-q)!}$.

f : $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$ by Theorem 1 and $F^p H^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ should be spanned by $\overline{\text{reg}}_{p,p-q} \text{CH}^p(\mathcal{X}_T, p-q)$ for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Conjecture 1; by Theorem 3 there is a variety Y such that $\overline{\text{reg}}_{p,p-q} \text{CH}^p(\mathcal{X}_T, p-q) \neq 0$ for $c \sim \frac{d^{N-q}}{(N-q)!}$ and by Conjecture there should be an indecomposable higher cycle in $\text{CH}^p(\mathcal{X}_T, p-q)$ with a non zero image under $\overline{\text{reg}}_{p,p-q}$ for $c \sim (p-q+1) \frac{d^{N-q}}{(N-q)!}$.

g : as in f; by Theorem 4 there is a variety Y and an indecomposable higher cycle in $\text{CH}^p(\mathcal{X}_T, p-q)$ with a non zero image under $\overline{\text{reg}}_{p,p-q}$ for $c \sim 2 \frac{d^{N-q}}{(N-q)!}$.

h : $F^p H^{p+q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) = 0$ for $c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$ by Theorem 1 and $F^p H^{p+q}(\mathcal{X}_T, \mathbb{C})_v$ should be spanned by $\overline{\text{reg}}_{p,p-q}(\text{CH}^p(\mathcal{X}_T, p-q))$ for $c \leq b \frac{d^{N-q}}{(N-q)!}$ by Conjecture 1. Here the situation is somewhat mysterious since I do not have any example of a non-zero element in $\text{CH}^p(\mathcal{X}_T, p-q)$.

1. PROOF OF THEOREM 1 AND OF ITS COROLLARY

1.1. Reduction to the local case. A straightforward adaptation of Nori's argument shows that if $H^{q+1}(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p) = 0$ for all $t \in T$ then $H^{q+1}(\mathcal{Y}_T, \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p) = 0$ ([No], Lemma 2.1). This shows that Theorem 1 is implied by the following purely local-analytic statement.

Proposition 1.2. *For all $\varepsilon \in \mathbb{R}_+^*$ there is an integer D depending only on p, ε, Y and on $\mathcal{O}(1)$ such that*

$$H^{q+1}(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p) = 0$$

for all integers $d \geq D$, $q \in \{-1, \dots, N-1\}$, for all smooth analytic varieties T endowed with a morphism $\phi: T \rightarrow H^0(Y, \mathcal{O}(d))$ such that

- for all $t \in T$ the cokernel of $d\phi(t)$ is of dimension at most $(1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$ and
- $\mathcal{X}_T$ is smooth.

1.3. Proof of proposition 1.2. Since the proof of proposition 1.2 proceeds by induction on several parameters, it is convenient to introduce the following notation. For a fixed positive integer d and a projective space $\mathbb{P}_{\mathbb{C}}^A$ I say that assertion $\mathcal{H}(Y, p, q, c, a)$ holds for the smooth projective variety $Y \subset \mathbb{P}_{\mathbb{C}}^A$ of dimension $N+1 > 0$ and for the integers $(p, q, c, a) \in \mathbb{N} \times \{-1, \dots, N-1\} \times \mathbb{N} \times \mathbb{Z}$ if

$$H^{q+1}(Y_t, \Omega_{Y_t, \mathcal{X}_T|Y_t}^p \otimes \mathcal{O}(a')) = 0$$

for all $a' \leq a$, for all smooth analytic varieties T endowed with a morphism $\phi: T \rightarrow H^0(Y, \mathcal{O}(d))$ such that $\mathcal{X}_T$ is smooth, and for all $t \in T$ such that the cokernel of $d\phi(t)$ is of dimension at most c .

The proof relies on the following four lemmas.

Recall (cf. [G1]) that for any positive integer c , a d -decomposition of c is the unique series $(c_d, \dots, c_r)$ such that $r \in \{1, \dots, d\}$, $c_d > \dots > c_r \geq r$ and

$$c = \binom{c_d}{d} + \dots + \binom{c_r}{r}.$$

Let

$$c_{<d>} = \binom{c_d - 1}{d} + \dots + \binom{c_r - 1}{r},$$

with the convention $\binom{\alpha}{\beta} = 0$ for $\alpha < \beta$.

The asymptotic behaviour of the function $c \mapsto c_{<d>}$ is described by the following lemma.

Lemma 1.3.1. *For all integers $M \geq 2$ there exists a function $f_M:]0, 1[\times \mathbb{N} \rightarrow]0, 1[$, $(\varepsilon, d) \mapsto \varepsilon'$ such that for all $\varepsilon \in]0, 1[$ $\varepsilon' \rightarrow \varepsilon^{\frac{M-1}{M}}$ for $d \rightarrow \infty$, and such that for all nonnegative integers*

$$c \leq (1 - \varepsilon) \frac{d^M}{M!}$$

one has

$$c_{<d>} \leq (1 - \varepsilon') \frac{d^{M-1}}{(M-1)!}.$$

The following lemma is the key ingredient of my proof. It relies on Green's hyperplane section theorem [G1].

Lemma 1.3.2. *Let Y' be a generic hyperplane section of Y . Then for $q \leq N-2$*

$$\left. \begin{array}{l} \mathcal{H}(Y', p, q, c_{<d>}, a) \\ \mathcal{H}(Y', p-1, q, c_{<d>}, a-1) \end{array} \right\} \Rightarrow \mathcal{H}(Y, p, q, c, a).$$

By convention, assumption $\mathcal{H}(Y', -1, q, c_{<d>}, a-1)$ holds.

The last two lemmas will only be used for $q = N-1$. Both rely on an idea of Green ([G2], p. 193-194).

Lemma 1.3.3. *Assume $c > 0$. Then*

$$\left. \begin{array}{l} \mathcal{H}(Y, p+1, q, c-1, a) \\ \mathcal{H}(Y, p+1, q-1, c, a) \end{array} \right\} \Rightarrow \mathcal{H}(Y, p, q, c, a).$$

By convention, assumption $\mathcal{H}(Y, p+1, -2, c, a)$ holds.

The last lemma has been proven by Nori for $a = 0$ ([No], remark 3.10); the general case involves no new ideas. See also [N] for computations of an explicit bound for s .

Lemma 1.3.4. *There is an integer s depending only on Y and $\mathcal{O}(1)$ such that for all $d \geq p + a + s$ the assertion $\mathcal{H}(Y, p, q, 0, a)$ holds for all $q \in \{-1, \dots, N-1\}$.*

Lemma 1.3.4 will only be used for $q = N-1$.

1.3.5. *Proof of proposition 1.2.* Postponing the proofs of the above Lemmas, I now show how they imply proposition 1.2.

I show more generally that for all $\varepsilon \in]0, 1[$ there is an integer D depending only on ε , p , a and Y such that assertion $\mathcal{H}(Y, p, q, c, a)$ holds for all $d \geq D$, $q \in \{-1, \dots, N-1\}$ and $c \leq (1 - \varepsilon) \frac{d^{N-q}}{(N-q)!}$. The case $a = 0$ is the proposition 1.2.

I proceed by induction on $N - q$, c and N .

Assume $N - q = 1$. The condition on c simplifies to $c \leq (1 - \varepsilon)d$. Choose D such that $\varepsilon D \geq p + a + s$. Thus for $d \geq D$ one has $d \geq (1 - \varepsilon)d + \varepsilon D \geq c + p + a + s$.

The case $c = 0$, $N - q = 1$ follows from Lemma 1.3.4.

The case $c > 0$, $N = 0$ and $q = -1$ follows from Lemma 1.3.3 by increasing induction on c . For the case $c > 0$, $N - q = 1$ let Y' be a generic hyperplane section of Y . Lemmas 1.3.2 and 1.3.3 provide successive implications

$$\left. \begin{array}{l} \mathcal{H}(Y, p+1, q, c-1, a) \\ \mathcal{H}(Y', p+1, q-1, c_{<d>}, a) \\ \mathcal{H}(Y', p, q-1, c_{<d>}, a-1) \end{array} \right\} \Rightarrow \mathcal{H}(Y, p+1, q-1, c, a) \Bigg\} \Rightarrow \mathcal{H}(Y, p, q, c, a).$$

Assertion $\mathcal{H}(Y, p, q, c, a)$ follows by increasing induction on both c and N .

Assume $N - q > 1$ and let Y' be a generic hyperplane section of Y . By induction on N and by Lemma 1.3.1 applied to $M = N - q$, assertions $\mathcal{H}(Y', p, q, c_{<d>}, a)$ and $\mathcal{H}(Y', p-1, q, c_{<d>}, a-1)$ hold. Hence by Lemma 1.3.2 assertion $\mathcal{H}(Y, p, q, c, a)$ holds. $\square$

1.4. Proof of the Lemmas.

1.4.1. *Proof of Lemma 1.3.1.* Let e be the smallest positive integer such that

$$(1 - \varepsilon) \frac{d^M}{M!} \leq \binom{M+d}{d} - \binom{M+e}{e}$$

for all $q \in \{0, \dots, N-1\}$. One has $e \sim \varepsilon^{1/M} d$ for $d \rightarrow \infty$.

The map $c \mapsto c_{<d>}$ is increasing. Thanks to the identities

$$\begin{aligned} \binom{M+d}{d} - \binom{M+e}{e} &= \sum_{i=1}^{d-e} \binom{M+d-i}{d-i+1} \\ \binom{M-1+d}{d} - \binom{M-1+e}{d-e} &= \sum_{i=1}^{d-e} \binom{M-1+d-i}{d-i+1} \end{aligned}$$

One gets

$$c_{<d>} \leq \binom{M-1+d}{d} - \binom{M-1+e}{e}.$$

The term on the right is asymptotically equivalent to $\varepsilon^{\frac{M-1}{M}} \frac{d^{M-1}}{(M-1)!}$ for $d \rightarrow \infty$. Define $\varepsilon' \in]0, 1[$ by $\varepsilon' = \varepsilon^{\frac{M-1}{M}}$ if $\frac{d^{M-1}}{(M-1)!} \geq \binom{M-1+d}{d} - \binom{M-1+e}{e}$ and by the formula

$$(1 - \varepsilon') \frac{d^{M-1}}{(M-1)!} = \binom{M-1+d}{d} - \binom{M-1+e}{e}$$

otherwise. Thus $\varepsilon' \in]0, 1[$ and $\varepsilon' \sim \varepsilon^{\frac{M-1}{M}}$ for $d \rightarrow \infty$. $\square$

1.4.2. *Proof of Lemma 1.3.2.* Let $i: Y' \rightarrow Y$ be the projective embedding, let $\mathcal{Y}'_T = Y' \times T$ and let $\mathcal{X}'_T = \mathcal{X}_T \times_{\mathcal{Y}_T} \mathcal{Y}'_T$. Fix $t \in T$.

Since Y' is generic, one may assume that $\mathcal{X}'_T$ is smooth and that X_t meets Y'_t transversally in Y_t . Restricting T if necessary, one may further assume that for all $u \in T$, X_u meets Y'_u transversally in Y_u . Hence there is a well-defined map $\phi': T \rightarrow \mathbb{P}H^0(Y', \mathcal{O}(d))$, constructed as follows.

Let $p_Y: H^0(Y, \mathcal{O}(d)) \setminus \{0\} \rightarrow \mathbb{P}H^0(Y, \mathcal{O}(d))$ be the natural projection and let

$$p_Y^* \phi: T \times_{\mathbb{P}H^0(Y, \mathcal{O}(d))} H^0(Y, \mathcal{O}(d)) \rightarrow H^0(Y, \mathcal{O}(d))$$

be the map deduced from ϕ by base change. Then ϕ' is defined by the map

$$p_{Y'}^* \phi': T \times_{\mathbb{P}H^0(Y, \mathcal{O}(d))} H^0(Y, \mathcal{O}(d)) \rightarrow H^0(Y', \mathcal{O}(d))$$

obtained by composing $p_Y^* \phi$ with the linear projection $H^0(Y, \mathcal{O}(d)) \rightarrow H^0(Y', \mathcal{O}(d))$ which maps to 0 the polynomials vanishing on Y' .

Let $E = p_Y^* \text{Im}(d\phi(t))$, *resp.* $E' = p_{Y'}^* \text{Im}(d\phi'(t))$. Let

$$\begin{aligned} c &= \dim \text{coker}(d\phi(t)) = \text{codim}(E, H^0(Y, \mathcal{O}(d))), \quad \text{resp.} \\ c' &= \dim \text{coker}(d\phi'(t)) = \text{codim}(E', H^0(Y', \mathcal{O}(d))). \end{aligned}$$

Let $S = \bigoplus_{\delta \in \mathbb{N}} S_\delta$, *resp.* $S' = \bigoplus_{\delta \in \mathbb{N}} S'_\delta$ denote the homogenous polynomial ring of $H^0(Y, \mathcal{O}(1))$, *resp.* $H^0(Y', \mathcal{O}(1))$. Let $I \subset S$, *resp.* $I' \subset S'$ denote the homogenous ideal of Y , *resp.* Y' . One has $c = \text{codim}(E + I_d, S_d)$, *resp.* $c' = \text{codim}(E' + I'_d, S'_d)$. Since $Y' \subset Y$ is generic, Green's hyperplane theorem [G1] asserts that

$$c' \leq c_{<d>}.$$

Hence the cokernel of $d\phi'(t)$ is of dimension at most $c_{<d>}$.

The sheaf $\Omega_{\mathcal{Y}_T, \mathcal{X}_T | \mathcal{Y}_T}^p$ fits into the short exact sequence

$$0 \longrightarrow \Omega_{\mathcal{Y}'_T, \mathcal{X}'_T}^{p-1} \otimes \mathcal{O}'_Y(-1) \xrightarrow{dH^\wedge} \Omega_{\mathcal{Y}_T, \mathcal{X}_T | \mathcal{Y}'_T}^p \longrightarrow \Omega_{\mathcal{Y}'_T, \mathcal{X}'_T}^p \longrightarrow 0$$

where $H \in H^0(Y, \mathcal{O}(1))$ is an equation of Y' . Restricting this sequence to Y'_t and tensoring it by $\mathcal{O}'_Y(a)$ one gets the associated long exact sequence

$$\begin{aligned} H^{q+1}(Y'_t, \Omega_{\mathcal{Y}'_T, \mathcal{X}'_T | Y'_t}^{p-1} \otimes \mathcal{O}'_Y(a-1)) &\xrightarrow{dH^\wedge} H^{q+1}(Y'_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T | Y'_t}^p \otimes \mathcal{O}'_Y(a)) \longrightarrow \\ &\longrightarrow H^{q+1}(Y'_t, \Omega_{\mathcal{Y}'_T, \mathcal{X}'_T | Y'_t}^p \otimes \mathcal{O}'_Y(a)). \end{aligned}$$

The term on the left vanishes by assumption $\mathcal{H}(Y', p-1, q, c_{<d>}, a-1)$ and the term on the right vanishes by assumption $\mathcal{H}(Y', p, q, c_{<d>}, a)$. Hence the middle term vanishes as well: one has

$$(1) \quad H^{q+1}(Y'_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y'_t}^p \otimes \mathcal{O}'_Y(a)) = 0.$$

For any coherent sheaf $\mathcal{F}$ on Y_t there is a short exact sequence

$$0 \longrightarrow \mathcal{F} \otimes \mathcal{O}(-1) \xrightarrow{H} \mathcal{F} \longrightarrow i_* i^* \mathcal{F} \longrightarrow 0,$$

Take $\mathcal{F} = \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p \otimes \mathcal{O}_Y(a)$ and consider the associated long exact sequence

$$\begin{aligned} H^{q+1}(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p \otimes \mathcal{O}_Y(a-1)) &\xrightarrow{H} H^{q+1}(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p \otimes \mathcal{O}_Y(a)) \\ &\longrightarrow H^{q+1}(Y'_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y'_t}^p \otimes \mathcal{O}'_Y(a)). \end{aligned}$$

The term on the right vanishes by (1). Since $q \in \{0, \dots, N-1\}$, by increasing induction on a the term on the right vanishes as well, the case $a \ll 0$ following from Serre's vanishing theorem. Hence the term in the middle vanishes. $\square$

1.4.3. Proof of Lemma 1.3.3. Since the assertion depends only on the tangent space of T at t , one may replace T by a small open neighbourhood of t in the tangent space of T at t ; hence one may assume $T = B \times S$, where $B \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ is an open neighbourhood of $\phi(t)$ of codimension at most c and S is an open ball.

Let $B' \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ be a smooth analytic variety containing B as a codimension 1 subvariety, and let $T' = B' \times S$. Then $\mathcal{Y}_T \subset \mathcal{Y}_{T'}$ is a smooth codimension 1 subvariety and one has the following short exact sequence of sheaves on $\mathcal{Y}_T$:

$$0 \longrightarrow \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^p \longrightarrow \Omega_{\mathcal{Y}_{T'}, \mathcal{X}_{T'}|_{\mathcal{Y}_T}}^{p+1} \longrightarrow \Omega_{\mathcal{Y}_T, \mathcal{X}_T}^{p+1} \longrightarrow 0.$$

Restrict this sequence to Y_t , tensorise it with $\mathcal{O}(a)$ and consider the associated long exact sequence

$$H^q(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^{p+1} \otimes \mathcal{O}(a)) \longrightarrow H^{q+1}(Y_t, \Omega_{\mathcal{Y}_T, \mathcal{X}_T|Y_t}^p \otimes \mathcal{O}(a)) \longrightarrow H^{q+1}(Y_t, \Omega_{\mathcal{Y}_{T'}, \mathcal{X}_{T'}|Y_t}^{p+1} \otimes \mathcal{O}(a)).$$

The terms on the right and on the left vanish by assumption, hence the term in the middle vanishes as well. $\square$

2. PROOF OF THEOREM 2

Consider the exact sequence

$$R^k \pi_* \mathbb{C} \rightarrow R^k \pi_*(j_{T*} \mathbb{C}) \rightarrow R^k \pi_*(\mathbb{C} \rightarrow j_{T*} \mathbb{C}) \rightarrow R^{k+1} \pi_* \mathbb{C} \rightarrow R^{k+1} \pi_*(j_{T*} \mathbb{C})$$

and let $\mathcal{H}_{\mathcal{X}_T, v}^k = \text{coker}(R^k \pi_* \mathbb{C} \rightarrow R^k \pi_*(j_{T*} \mathbb{C}))$ denote the locally trivial subsheaf of $R^k \pi_{Y*}(\mathbb{C} \rightarrow j_{T*} \mathbb{C})$ whose fiber at any point $t \in T$ is canonically isomorphic to $H^k(X_t, \mathbb{C})_v$. Let $F^p H^0(T, \mathcal{H}_{\mathcal{X}_T, v}^k)$ denote the space of sections of $\mathcal{H}_{\mathcal{X}_T, v}^k$ whose fiber at any point $t \in T$ belongs to $F^p H^k(X_t, \mathbb{C})_v$.

In [O] I have shown

Theorem ([O]). *For all integers $b \in \mathbb{N}^*$ there is an integer D depending only on b , Y and on $\mathcal{O}(1)$, such that for all $d \geq D$, for all $q \in \{[(N+1)/2], \dots, N-1\}$, for all*

$$c \leq b \frac{d^{N-q}}{(N-q)!}$$

for all codimension c algebraic varieties $T \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ lying in the locus parametrizing smooth hypersurfaces and for all $t \in T$, the following holds.

If N is even and $q = N/2$ then the space $F^{N-q}H^0(T, \mathcal{H}_{\mathcal{X}_T, v}^N)$ is spanned by families of algebraic cycles of codimension q and degree at most b .

If $q > N/2$ then $F^{N-q}H^0(T, \mathcal{H}_{\mathcal{X}_T, v}^N) = 0$.

The proof of [O] gives in fact a stronger result, namely one can replace the subspace $T \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ by any algebraic variety T endowed with a morphism $\phi: T \rightarrow \mathbb{P}H^0(Y, \mathcal{O}(d))$, such that

- $\phi(T)$ lies in the locus parametrizing smooth hypersurfaces and
- for all $t \in T$ the cokernel of the differential $d\phi(t): T_{T,t} \rightarrow T_{\mathbb{P}H^0(Y, \mathcal{O}(d)), \phi(t)}$ is of dimension at most c ;

This stronger version of the theorem of [O] is equivalent to Theorem 2. Indeed, one only needs to show

$$F^{N-q}H^N(\mathcal{X}_T, \mathbb{C})_v \simeq F^{N-q}H^0(T, \mathcal{H}_{\mathcal{X}_T, v}^N).$$

But the Lefschetz hyperplane Theorem and the Leray spectral sequence give a canonical isomorphism

$$H^N(\mathcal{X}_T, \mathbb{C})_v \simeq H^0(T, \mathcal{H}_{\mathcal{X}_T, v}^N).$$

Since for any $t \in T$, the restriction map $H^N(\mathcal{X}_T, \mathbb{C})_v \rightarrow H^N(\mathcal{X}_t, \mathbb{C})_v$ is a homomorphism of Hodge structures, hence is strict, the result follows. $\square$

3. PROOF OF THEOREM 3

Since $q \geq N/2$, the generic hypersurface of degree d containing V is smooth; let $T \subset \mathbb{P}H^0(Y, \mathcal{O}(d))$ be a smooth open affine subvariety of the space of all smooth hypersurfaces containing V . One has $c \sim_{d \rightarrow \infty} \frac{d^{N-q}}{(N-q)!}$.

For all $i \in \{1, \dots, p-q\}$ choose hyperplane sections $H_i \subset T$ in general position and let $f_i \in H^0(T, \mathcal{O}_T)$ be the equations of H_i ; Let $P_i = \bigcap_{j>i} H_j$, $Q_i = \bigcup_{j \leq i} H_j$ and $T_i = P_i \setminus (P_i \cap Q_i)$. For $j \leq i$ the functions f_j do not vanish on T_i , hence belong to $H^0(T_i, \mathcal{O}_{T_i}^*)$ and there are well-defined higher cycles in $\text{CH}^{q+i}(\mathcal{X}_{T_i}, i)$

$$Z_i = \psi_{1,1}^i([V \times T_i], f_1, \dots, f_i).$$

I show by increasing induction on i that $\overline{\text{reg}}_{q+i,i}(Z_i) \neq 0$. Case $i = p-q$ implies Theorem 3.

Case $i = 0$ a straightforward adaptation of [V2], Proposition 4: for $d \gg 0$ and for any open subset $U_0 \subset T_0$ the cohomology class of the cycle $[V \times U_0] \in \text{CH}^q(\mathcal{X}_{U_0})$ in $H^{2q}(\mathcal{X}_{U_0}, \mathbb{C})$ does not belong to $j_{U_0}^*(H^{2q}(\mathcal{Y}_{U_0}, \mathbb{C}))$.

Assume $i > 0$. Then $T_{i-1} \cup T_i = P_i \setminus (P_i \cap Q_{i-1})$, hence T_{i-1} is a hyperplane section of $T_{i-1} \cup T_i$ and T_i is the complementary open set. Hence there is a commutative diagram

$$\begin{array}{ccc} \mathrm{CH}^{q+i}(\mathcal{X}_{T_i}, i) & \xrightarrow{l_{q+i,i}} & \mathrm{CH}^{q+i-1}(\mathcal{X}_{T_{i-1}}, i-1) \\ \downarrow \overline{\mathrm{reg}}_{q+i,i} & & \downarrow \overline{\mathrm{reg}}_{q+i-1,i-1} \\ F^{q+i} \mathrm{H}^{2q+i+1}(\mathcal{Y}_{T_i}, \mathcal{X}_{T_i}, \mathbb{C}) & \xrightarrow{\mathrm{res}} & F^{q+i-1} \mathrm{H}^{2q+i}(\mathcal{Y}_{T_i}, \mathcal{X}_{T_{i-1}}, \mathbb{C}) \end{array}$$

where res is the residue map and $l_{q+i,i}$ is the linking map for the long exact sequence of higher Chow groups. One has

$$l_{q+i,i}(Z_i) = (l_{q,0} \circ \psi_{1,1}^i)([V \times T_i], f_1, \dots, f_i) = \psi_{1,1}^{i-1}([V \times T_{i-1}], f_1, \dots, f_{i-1}) = Z_{i-1}.$$

The induction assumption $\overline{\mathrm{reg}}_{q+i-1,i-1}(Z_{i-1}) \neq 0$, and the commutativity of the diagram implies $\overline{\mathrm{reg}}_{q+i,i}(Z_i) \neq 0$. $\square$

4. PROOF OF THEOREM 4

4.1. Construction of the cycle Z_U . Let $P \subset Y$ be a codimension q linear subspace, let $R \subset Y$ be a codimension $q+2$ linear subspace such that $R \subset P$ and let S be the blow-up of P along R . There are natural maps $\psi: S \rightarrow \mathbb{P}_{\mathbb{C}}^1$ (where $\mathbb{P}_{\mathbb{C}}^1$ parametrizes hyperplanes of P containing R) and $s: S \rightarrow Y$. For any $x \in \mathbb{P}_{\mathbb{C}}^1$ let $L_x = s(\psi^{-1}(x))$ denote the codimension $q+1$ linear subspace of Y . For any subvariety $B \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ let $C_B = S \times_Y \mathcal{X}_B$ and let $\psi_B: C_B \rightarrow \mathbb{P}_{\mathbb{C}}^1$ and $s_B: C_B \rightarrow \mathcal{X}_B$ be the morphisms deduced from ψ and s by base change.

Let $U \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ be the locus of hypersurfaces X_t such that $[L_0 \cap X_t] = dR$ and $[L_{\infty} \cap X_t] = dR$. One has $\mathrm{codim}(U, \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))) \sim_{d \rightarrow \infty} 2 \frac{d^{N-q}}{(N-q)!}$. By construction, one has $s_U(\psi_U^{-1}(0)) = R \times U = s_U(\psi_U^{-1}(\infty))$, hence the couple (C_U, ψ_U) defines an element $Z_U \in \mathrm{CH}^{q+1}(U, 1)$.

4.2. Proof of $\overline{\mathrm{reg}}_{q+1,1}(Z_U) \neq 0$. Let $T \subset \mathbb{P}\mathrm{H}^0(Y, \mathcal{O}(d))$ be the locus of hypersurfaces X_t such that $L_0 \subset X_t$ and $[L_{\infty} \cap X_t] = dR$. Then $T \subset T \cup U$ is a hyperplane section and $U \subset T \cup U$ is the complementary open set. Hence there is a natural linking homomorphism $l_{q+1,1}: \mathrm{CH}^{q+1}(\mathcal{X}_U, 1) \rightarrow \mathrm{CH}^q(\mathcal{X}_T)$. Since C_T is the union of a variety dominating $\mathbb{P}_{\mathbb{C}}^1$ and of the variety $\psi_T^{-1}(0) \simeq L_0 \times T$, one has $l_{q+1,1}(Z_U) = d[s_T(\psi_T^{-1}(0))] = d[L_0 \times T]$.

I now proceed as in section 3. A straightforward adaptation of the argument of [V2], Proposition 3 shows that for $d \gg 0$ the class $\mathrm{cl}_q(d[L_0 \times T]) \in \mathrm{H}^{2q}(\mathcal{X}_T, \mathbb{C})$ does not belong to $j_T^*(\mathrm{H}^{2q}(\mathcal{Y}_T, \mathbb{C}))$, hence $\overline{\mathrm{cl}}_q(d[L_0 \times T]) \in \mathrm{H}^{2q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C})$ is non-zero. Since the diagram

$$\begin{array}{ccc} \mathrm{CH}^{q+1}(\mathcal{X}_U, 1) & \xrightarrow{l_{q+1,1}} & \mathrm{CH}^q(\mathcal{X}_T) \\ \downarrow \overline{\mathrm{reg}}_{q+1,1} & & \downarrow \overline{\mathrm{cl}}_q \\ \mathrm{H}^{2q+2}(\mathcal{Y}_U, \mathcal{X}_U, \mathbb{C}) & \xrightarrow{\mathrm{res}} & \mathrm{H}^{2q+1}(\mathcal{Y}_T, \mathcal{X}_T, \mathbb{C}) \end{array}$$

is commutative, it follows that $\overline{\mathrm{reg}}_{q+1,1}(Z_U) \neq 0$.

4.3. Indecomposability of the cycle Z_U . Since $\overline{\text{reg}}_{q+1,1}(Z_U) \neq 0$, it is enough to show that the space of decomposable cycles is contained in the kernel of $\overline{\text{reg}}_{q+1,1}$.

Let $i \in \{1, \dots, q\}$, $Z'_U \in \text{CH}^i(\mathcal{X}_U)$ and $Z''_U \in \text{CH}^{q-i}(\mathcal{X}_U, 1)$. One has to show $\text{reg}_{q+1,1}(\phi_{i,0}(Z'_U \otimes Z''_U)) \in j_U^* F^{q+1} H^{2q+1}(\mathcal{Y}_U, \mathbb{C})$. Since the diagram

$$\begin{array}{ccc} \text{CH}^i(\mathcal{X}_U) \otimes \text{CH}^{q+1-i}(\mathcal{X}_U, 1) & \xrightarrow{\phi_{i,0}} & \text{CH}^{q+1}(\mathcal{X}_U, 1) \\ \downarrow \text{cl}_i \otimes \text{reg}_{q+1-i,1} & & \downarrow \text{reg}_{q+1,1} \\ F^i H^{2i}(\mathcal{X}_U, \mathbb{C}) \otimes F^{q-i+1} H^{2q-2i+1}(\mathcal{X}_U, \mathbb{C}) & \xrightarrow{\sim} & F^{q+1} H^{2q+1}(\mathcal{X}_U, \mathbb{C}), \end{array}$$

is commutative, it is enough to show $\text{cl}_i(Z'_U) \in H^{2i}(\mathcal{Y}_U, \mathbb{C})$ and $\text{reg}_{q+1-i,1}(Z''_U) \in H^{2q-2i+1}(\mathcal{Y}_U, \mathbb{C})$.

If $i < q$ this follows from the isomorphisms given by Theorem 1,

$$\begin{aligned} F^i H^{2i}(\mathcal{X}_U, \mathbb{C}) &\simeq F^i H^{2i}(\mathcal{Y}_U, \mathbb{C}), \\ F^{q-i+1} H^{2q-2i+1}(\mathcal{X}_U, \mathbb{C}) &\simeq F^{q-i+1} H^{2q-2i+1}(\mathcal{Y}_U, \mathbb{C}). \end{aligned}$$

If $i = q$, one has

$$\text{CH}^1(\mathcal{X}_U, 1) \simeq H^0(U, \mathcal{O}^*(U)) \simeq \text{CH}^1(\mathcal{Y}_U, 1),$$

hence $\text{reg}_{1,1}(Z''_U) \in H^1(\mathcal{Y}_U, \mathbb{C})$; on the other hand, since $\mathcal{X}_U$ does not contain any flat family of cycles of codimension q and degree less than two, by Theorem 2 one has $F^q H^{2q}(\mathcal{X}_U, \mathbb{C}) \simeq F^q H^{2q}(\mathcal{Y}_U, \mathbb{C})$, hence $\text{cl}_q(Z'_U) \in H^{2q}(\mathcal{Y}_U, \mathbb{C})$. $\square$

4.4. Indecomposability of the restriction of Z_U to a very general fiber $\mathcal{X}_u$, $u \in U$. This follows from the indecomposability of Z_U by Proposition 5 of [V2].

Acknowledgments. I would like to thank Stefan Müller-Stach for explaining to me what a higher Chow group is, and for a useful remark on conjecture 1.

REFERENCES

- [B] S. BLOCH *Algebraic cycles and higher K-theory*. Adv. in Math. 61, no. 3, p. 267–304. (1986).
- [G1] M. GREEN. *Restrictions of linear series to hyperplanes, and some results of Macaulay and Gotzmann*. Algebraic curves and projective geometry, p. 76–86, LNM 1389, (1989).
- [G2] M. GREEN. *Koszul cohomology and geometry*. Lectures on Riemann surfaces (Trieste, 1987), p. 177–200, Algebraic Curves and Projective Geometry (E. Ballico and C. Ciliberto, eds), p. 77–88, Springer LNM 1389, Berlin (1989).
- [N] J. NAGEL. *Effective bounds for Hodge-theoretic connectivity*. J. Alg. Geom. 11, no 1, p. 1–32, (2002).
- [No] M. NORI. *Algebraic cycles and Hodge theoretic connectivity*. Invent. math. 111 (2), p. 349–373 (1993).
- [O] A. OTWINOWSKA *Variétés de Hodge*. Submitted, (2001).
- [V2] C. VOISIN. *Nori's connectivity theorem and Higher Chow groups*. J. Inst. Mat. Jussieu, no 2, p. 307–329 (2000).